



# TruRange™



# Product

## information sheet

## Innovative and Sustainable Pasture Management

### // BEST USES

TruRange™ provides extended control of over 70 weed and brush species including Canada thistle, dandelion, leafy spurge, western snowberry and wild rose for up to 24 months.

### // KEY STRENGTHS

TruRange offers a new way for ranchers to provide high quality forage for their cattle in an environmentally sustainable and effective manner. TruRange provides residual weed and brush control in one easy-to-use, low-rate product with no grazing restrictions. With less active ingredient going into the environment, TruRange brings innovation and sustainability to pasture and rangeland management.

#### *TruRange Rates*

##### **Weed Rate (68 g/acre or 167 g/ha)**

- 1 jug (1.361 kg) treats 20 acres

##### **Brush Rate (135 g/acre or 334 g/ha)**

- 1 jug (1.361 kg) treats 10 acres

#### *Efficacy*

- TruRange delivers extended broad-spectrum control of broadleaf weeds and encroaching brush
- It promotes the grass understory and helps preserve desirable vegetation
- Two modes of action for effective resistance management
- Quickly taken up by the leaves, stems and roots of plants
- Rain-fast at four hours after application

#### *Grazing and Haying Restrictions*

- There are no grazing or haying restrictions for non-lactating or lactating animals (including cattle, horses, sheep and goats) when using TruRange herbicide as directed
- Grazing animals do not have to be moved off the rangeland or pasture before, during or after applying TruRange herbicide
- See Manure Management and Hay and Other Plant Material Management on next page for additional information

---

#### // Active Ingredients

Metsulfuron-methyl  
12.6%  
Aminocyclopyrachlor  
39.5%

#### // Mode of Action

Auxinic and  
ALS inhibitor

#### // Group

2 and 4

#### // Formulation

Wettable granule

#### // Packaging

Case = 8 x 1.361 kg

## Manure Management

Aminocyclopyrachlor, an ingredient in TruRange herbicide, passes through an animal's digestive tract and is excreted in urine and manure at levels that may cause injury to susceptible plants.

- Do not transfer grazed animals from areas treated with TruRange herbicide to areas where sensitive crops occur without first allowing three days of grazing on untreated areas

The following restrictions apply to manure from animals that have grazed forage or eaten hay from areas that have been treated with TruRange herbicide within the prior 18 months:

- Do not apply manure to land used for growing susceptible crops
- Manure may only be applied on rangeland
- Do not use manure as mulch or compost and do not apply directly on or around desirable plants
- Manure must only be used on-farm

After removing animals from grazing on treated areas or eating forage or hay from treated areas, and waiting three days for treated material to clear the animal's digestive system, the animal's manure is no longer subject to the above restrictions.

## Hay and Other Plant Material Management

The following restrictions apply to all plant materials from areas treated with TruRange herbicide within the prior 18 months:

- Do not use plant material as mulch or compost and do not apply directly on or around desirable plants
- Hay cut from grass that has been treated with TruRange herbicide within the prior 18 months must only be used on-farm
- Plant material from the treated area is no longer subject to the above restrictions 18 months after treatment

## Application Information

**Ground application:** 20 gal/ac of water volume (200 L/ha) minimum

**Aerial application:** 3-5 gal/ac of water volume (30-50 L/ha) minimum



Darrell Chambers  
British Columbia  
403-498-7006  
darrell.chambers@bayer.com

Megan Coverdale  
Alberta  
587-434-3411  
megan.coverdale@bayer.com

Brett Armstrong  
Saskatchewan and Manitoba  
403-461-6008  
brett.armstrong@bayer.com

Will Roberts  
Ontario and Atlantic Canada  
226-821-5845  
will.roberts@bayer.com

es.bayer.ca 1-888-283-6847