



Bayer Greenhouse Vegetable Production Solutions

PRODUCTS	ACTIVE	GROUP	FAMILY OR SUBGROUP	SYSTEMIC OR CONTACT	MODE OF ACTION	KEY PESTS	CROP USES	RESIDUAL CONTROL	REI	PHI (DAYS)
INSECT AND MITE SOLUTIONS										
Altus™	flupyradifurone	4D	butenolide	Systemic/contact xylem mobile	agonist of the nicotinic acetylcholine receptor rapid inhibition of feeding	aphids, whitefly, leafhoppers	tomato, cucumber*, lettuce, pepper, strawberries	Foliar 14-21 days Drench up to 6 weeks	12hr	12 hours – strawberries 1 day – tomato, cucumber, lettuce 3 days – peppers, eggplant
Kontos™	spirotetramat	23	tetramic acid	Fully systemic – xylem and phloem mobile	lipid biosynthesis inhibitor	aphids, whitefly	tomato, cucumber, pepper, eggplant, lettuce	Up to 30 days	12hr	3 days 7 days – lettuce
Forbid™	spiromesifen	23	tetramic acid	Contact local systemic – translaminar	lipid biosynthesis inhibitor	mites, whitefly	tomato, cucumber, pepper, eggplant	Up to 30 days	12hr	3 days
Intercept™	imidacloprid	4	neonicotinoid	Systemic/contact – xylem mobile	binds to post synaptic area nerve	aphids, whitefly, swede midge	tomato, cucumber, pepper, eggplant, head and stem brassica (Group 5a) – see updated label for details	Up to 8 weeks		3 days – pepper, eggplant 1 day – cucumber, tomato 21 days – brassica 28 days – lettuce
DISEASE CONTROL SOLUTIONS										
Rhapsody™ ASO	bacillus subtilis – QST 713	44	biologic	Contact	multi site – lipopeptides puncture membranes, antibacterial action, triggers plants internal defenses	powdery mildew, botrytis, downy mildew, sclerotinia rot, gummy stem blight, cercospora leaf spot, white mould, bacterial blight, bacterial speck, bacterial spot, pin rot	tomato, cucumber, pepper, eggplant, lettuce, leafy vegetables, cole crops, cucurbits, herbs and spices Refer to label for specific crops and labelled pests	7-10 days, except leafy vegetables (10-14 days)	4 hr or until dry	0 days
Previcur™ N	propamocarb	28	carbamate	Systemic	multi site – reduces spores and mycelium, inhibits membrane function	pythium, downy mildew (suppression) cucumber only	tomato, cucumber, pepper, eggplant	7-10 days – pythium 7-14 days – downy mildew	12hr	1 day – tomato, peppers, eggplants 2 days – cucumber
Scala™	pyrimethanil	9	analinopyrimidine	Local systemic, vapour activity	single site – inhibits enzyme secretion of fungus	botrytis	tomato, cucumber	14 days	12hr	1 day
Luna Privilege™	fluopyram	7	SDHI (succinate de-hydrogenase inhibitor)	Systemic – xylem mobile	single site – C2 Respiration – inhibition of complex II	botrytis, powdery mildew, early blight	cucumber, pepper, eggplant, tomato, lettuce	Foliar 7-14 days Drench 30 days	12hr	Can be applied up until the day of harvest for tomato, pepper and cucumber 7 days – lettuce

*May cause some leaf yellowing and/or mottling.

ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS.

Altus™, Forbid®, Intercept™, Kontos™, Luna Privilege™, Previcur™, Rhapsody®, Scala®, is a trademark of Bayer Group. Used under license. Bayer CropScience Inc. is a member of CropLife Canada. ©2021 Bayer Group. All rights reserved.